

()

//

x

(T4 T3)

(p< /)

(P> /)

(p< /)

(T4 T3)

(P> /)

x

.()

.()
.()

.()

.()

x

)

)

(

(

x x /

.()

.()

.()

²UFFDA

(^ξ +^r)

(NSP)

2 . User-friendly feed formulation done agine

3 . Beta glucanase

4 . Beta xylanase

.()

1 . Non-starch Polysaccharides

(×)

(p < /)
%

(³T₃,⁴T₄)

()

()

(.)

()

()

()

SAS

(P > /)

$$Y_{ijk} = \mu + A_i + B_j + (AB)_{ij} + \beta X_{ij} + \varepsilon_{ij}$$

A_i μ Y_{ijk}
 $(AB)_{ij}$ B_j
 ε_{ijk} βX_{ij}

(p < /)

()

()

-
- 1 . Hen Day
 - 2 . Egg Mass
 - 3 . Thriiodothyronine
 - 4 . Thetraiodothyronine

...

:

/ a	/ a	/ a	/ a	%
± /	± /	± /	± /	
/ ab	/ ab	/ ab	/ a	%
± /	± /	± /	± /	
/ ab	/ ab	/ ab	/ a	%
± /	± /	± /	± /	
/ b	/ b	/ b	/ b	%
± /	± /	± /	± /	
/	/	/	/	
± /	± /	± /	± /	
/	/	/	/	
± /	± /	± /	± /	

(ng/ml)	(ng/ml)			
/ a	/ a	/	/	%
± /	± /	± /	± /	
/ ab	/ a	/	/	%
± /	± /	± /	± /	
/ b	/ b	/	/	%
± /	± /	± /	± /	
/ b	/ b	/	/	%
± /	± /	± /	± /	
/	/	/	/	
± /	± /	± /	± /	
/	/	/	/	
± /	± /	± /	± /	

(SE)

(p < /)

a-b

(ng/ml)		(ng/ml)			
a /	a /	a /	a /	a /	a /
± /	± /	± /	± /	± /	± /
ab /	ab /	b /	a /	a /	a /
± /	± /	± /	± /	± /	± /
ab /	b /	bc /	b /	bc /	bc /
± /	± /	± /	± /	± /	± /
/ b	/ b	/ c	/ b	/ c	/ c
± /	± /	± /	± /	± /	± /
/	/	/	/	/	/
± /	± /	± /	± /	± /	± /
/	/	/	/	/	/
± /	± /	± /	± /	± /	± /

(p < /)

a b

(SE)

()

(p < /)

()

(P > /)

%

NSP

()

(p < /)

(P > /)

(p < /)

()

()

()

()

()

REFERENCES

(Brassica Napus)

3. Anonymous. 2004. L. I. R. Parent stock management program. Cuxhven, Germany.
4. Bao, C., F. Xu., Y. Sun., C. Shao., & M. Liu. 1995. Histological changes in main viscera of broiler chicks fed on different rapeseed meals containing toxic substances. *Jiangsu Journal Agricultural Science*. 11: (4) 34-40.
5. Bedford, M. R. 2000. Exogenous enzyme in monogastric nutrition, their current value and future benefits. *Animal Feed Science and Technology*. 86:1-13.
6. Classen, H. L. & G. L. Campbell. 1991. The use of enzyme in diets for laying hens. *Proceedings of the Expert Committee on Animal Nutrition*, June 3-4, Edmonton, AB.
7. Friedrich, S., L. Matthias., H. Horst., J. Gerhard. & T. Frank. 2001. Rapeseed glucosinolates and iodine in sows affect the milk iodine concentration and the iodine status of piglets. *British Journal of Nutrition*, 85, 659-670.
8. Guenter, W., B. A. Slominski. & L. D. Campbell. 2001. Enhancement the feeding value of canola meals using exogenous enzymes. 12th Project Report, Canola Council of Canada, Canada. 45-49.

9. Hill, R. 1979. A review of the toxic effects of rapeseed meal whit observation on meal from improved varieties. *British Veterinary Journal*. 135: 3-16.
10. Huthail, N. & A. Suliaman. 2004. The effect of incorporating different levels of locally produced canola seeds (*Brassica napus, l.*) in the diet of laying hen. *International Journal of Poultry Science* 3(7): 490-496.
11. Kiiskinen, T. 1989. Effect of long-term use of rapeseed meal on egg production. *Annales Agriculture Fenniae*. 28:385-396.
12. Kohler H, A. Taurog., & H. B. Dunford. 1988. Spectral studies with lactoperoxidase and thyroid peroxidase: Interconversions between native enzyme, Compound II, and Compound III. *Archive of Biochemistry and Biophysics*, 264. 438-439.
13. Leeson, S. & J. D. Summers. 1991. Commercial poultry nutrition. Guelph, Ontario, Canada.
14. PAIK, I.K. & A.R. Robblee. 1984. Influence of feeding canola meal and seed on the level of thiocyanate ions in blood plasma and eggs of laying chickens. *Han'guk Ch' uksan Hakhiechi*, 26, 541-544.
15. Rotkiewicz, T. M. A. Kozłowski., S. M. Faruga., D. Wisniewska., H. Ikulski., D. Kozłowska., K. Rotiewicz and K. Kozłowski. 1993. Pathomorphological examination of thyroid glands and livers of broiler chickens fed with diets with maximum content of extracted rapeseed oil meal of the Polish Jantar cultivar and Canadian canola. *Acta-Academiae-Agriculturae-ac-Technicae-Olstenensis-Veterinaria*. 21, 219-228.
16. Rotter, B. A., O. D. Friesen., W. Guenter & R. R. Marquardt. 1990. Influence of enzyme supplementation on the bioavailable energy of barley. *Poultry Science*. 69: 1174-1181.
17. Sandman, M. & W. J. Schon. 1987. The proteins in rapeseed. Coposition and characterization. *Proceeding 7th Intern. Conference Rapeseed*. Poznan. Poland.
18. Seth, J., F. J. Rutherford, & I. Mckenize. 1975. Solid-phase radioimmunoassay of thyroxine in untreated serum. *Clin. Chem*. 21: 1406-1413.
19. Smits, C. H. M. & G. Annison. 1996. Non-starch poly saccharides in broiler nutrition-towards a physiologically valid approach to their determination. *World's Poultry Science Journal*. 52: 203-221.
20. Summers, J. D., S. Leeson & D. Spratt. 1988. Canola meal and egg size. *Canadian Journal of Animal Science*. 68: 907-913.
21. Zeb, A. Stter & U. Meulen. 1999. Effect of feeding different levels of rapeseed meal on the performance of broiler chickens. *Archive Fure Gfluegelkund*. 63: 77-81.