

( )

**(*Arachis hypogaea* L.)**

\*

( // : // : )

)

)

(

(

/

:

(Williams et al.,

.1995)

(Anis et al., 2001;

(Anis et al., 2001;

.Gahaner & Ashir, 1974)

.Arnon, 1989; Bell et al., 1987)

(Gahaner & Ashir,

.1974; Hausar et al., 1979; Widaryanto, 1994)

.(Craufurrd et al., 2002)

(Gahaner & Ashir, 1974;  
Hausar et al., 1979; Widaryanto, 1994; Yilmas,  
.1999)

(Kirby &  
Kitbamroon, 1986; Koochaki & Khalghani, 1996;  
.Mozing & Coffelt, 1984)

.(Choudhury et al., 1997)

.(Munda & Patel, 1989)

(Ghosh &  
.Biswas, 1995; Kirby & Kitvbmroong, 1986)

.(Gahaner & Ashir, 1974)

(Choudhury et al., 1997; David et al.,  
.2001)

(Gardner & Auma, 1989; Hanna et al.,  
.1994)

(Choy et al., 1982; Hausar et al., 1979; Hauser &  
Buchanan, 1981)  
(Kvin & Bergmark, 1987;  
.Mixon, 1989; Widaryanto, 1994)

.(Jaffar & Gardner, 1988)

.(Choudhury et al., 1997)

.(Ghosh & Biswas, 1995)

(Craufurrd et al., 2002)

(Hauser & Buchanan, 1981)

(Mozing & Coffelt, 1984)

(Model Li\_Cor 3100)

pH (

(1995) Williams et al.

( x / )+

( )

/ x / x

x x x

( x ) x

)

(

/ x / / x /

x

/( )

$$= \frac{(\quad \times /)}{\quad}$$

$$= \frac{\quad}{\quad}$$

$$= \frac{\quad}{\quad}$$


/ / /  
/( )  
/

SAS


MSTATC

EXCEL

/( )



/( )



/



/ /

...

:

| (meq/100gr) | (meq/100gr) | (meq/100gr) | (mg/kg) | (mg/kg) | (mg/kg) | (mg/kg) |
|-------------|-------------|-------------|---------|---------|---------|---------|
| /           | /           | /           | /       | /       | /       | /       |
| / *         | / ns        | / ns        | / *     | / ns    | / ns    | / ns    |
| / **        | / ns        | / *         | / **    | / *     | / **    | / *     |
| / **        | / *         | / *         | / **    | / **    | / **    | / **    |
| / ns        | / ns        | / ns        | / ns    | / ns    | / ns    | / ns    |
| /           | /           | /           | /       | /       | /       | /       |
| /           | /           | /           | /       | /       | /       | (%)     |

\* \*\*, ns






( )


/

/


/



تراکم کاشت (بوته در متر مربع)




( )  
( )




( )  
( )  
( )  
( )

...  
:  
(Gahaner & Ashir, 1974; Choudhury et al., 1997; Kirby & Kitbamroong, 1986)



(Gardner & Auma, 1989; Koochaki & Khalghani, 1996; Widaryanto, 1994)



(Choudhury et al., 1997; David et al., 2001; Hanna et al., 1994)

(Gahaner & Ashir, 1974; Choudhury et al., 1997; Jaffar & Gardner, 1988; Yilmas, 1999)

(Wynne, 1974)

(Gahaner & Ashir, 1974)

(Gardner & Auma,  
1989; Kirby & Kitbamroong, 1986; Kvin &  
Bergmark, 1987; Munda & Patel, 1989)

/

/

/

/

(Choy et al., 1982; Craufurrd et al., 2002;  
.Ghosh & Biswas, 1995)

/

/

/

(Gahaner & Ashir, 1974;  
Choudhury et al., 1997; Koochaki & Khalghani,  
1996; Mozing & Coffelt, 1984)

/

(Gahaner

.& Ashir, 1974; Hanna et al., 1994)

/

)

(

/



## REFERENCES

1. Anis, R., Wells, R. & Thomas, G. (2001). Reproductive allocation on branching of Virginia- type peanut cultivars bred for yield in North Carolina. *Crop Sci*, 41, 72- 77.
2. Arnon, I. (1989). *Processing tropical crops*. Macinillan press LTD. PP. 124- 140.
3. Bell, M. J., Muchow, R. C. & Wilson, G. L. (1987). The effect of plant population on peanut (*Arachis hypogaea* L.) in a monsoonal tropical environment. *Field Crop Research*, 17, 91- 107.
4. Gahaner, A. & Ashir, A. (1974). Vegetative and reproductive development of virginia-type peanut varieties in different stand densities. *Crop Sci*, 14, 412- 416.
5. Choudhury, B., Zahid, M., Ghosh, P. & Samui, R. C. (1997). Effect of rate and row spacing on confectionary groundnut in West Bengal, India. *International Arachis Newsletter*, 17, 71- 73.
6. Choy, E. W., Ston, C. J. F., Matlock, R. S., Mccauley, G. N. & Chin- Choy, E. W. (1982). Plant population and irrigation effects on spanish peanuts. *Peanut Sci*, 2, 73- 76.
7. Craufurrd, P. Q., Vara, P. V. & Summerfield, R. J. (2002). Dry matter production and rate of change of harvest index at high temprature in peanut. *Crop Sci*, 42, 146-151.
8. David, L., Jordan, B., Beam, P., Dewayne, J. & Spears, F. (2001). Peanut response to prohexadione calcium in three seeding rate – planting pattern system. *Agron J*, 93, 232- 236.
9. Gardner, F. P. & Auma, E. Q. (1989). Canopy structure, light interception and yield market quality of peanut genotypes as influenced by planting pattern and planting date. *Field Crop Research*, 20(1), 13- 29.
10. Ghosh, A. K. & Biswas, A. K. (1995). Regulation of correlative senescence in peanut (*Arachis hypogaea* L.) by source- sink alteration through physical and hormonal means. *Agron J*, 175, 195- 202.
11. Hanna, F. R., Salama, N. F. & Gawad, M. (1994). Effect of population density on yield of peanut. *Annals of Agricultural Science moshtohor*, 32, 731- 742.
12. Hausar, E. W., Buchanun, G. A. & Slaughter, J. W. (1979). Effects of row spacing, weed- free maintenance periods and herbicide systems on the yield of florunner peanuts. In: *Proceedings of American Peanut Research and Education Society*. 11: 1- 4.
13. Hauser, E. W. & Buchanan, G. A. (1981). Influence of row spacing, seeding rate and herbicide systems on the competition and yield of peanut. *Peanut Sci*, 8(1), 74- 81.
14. Jaffar, Z. & Gardner, F. P. (1988). Canopy development yield and market quality in peanut as affected by genotype and planting pattern. *Crop Sci*, 28, 299- 305.
15. Kirby, J. S. & Kitbamroong, C. (1986). Growth and development of the runner Virginia market type peanut (*Arachis hypogaea* L.). In: *Proceedings of American Peanut Research and Education Society*. 18: 48.
16. Koochaki, A. & Khalghani, J. (1996). *Understanding Crop Production*. Fersowsi University of Mashhad Press. 260p. (In Farsi).
17. Kvin, C. S. & Bergmark, C. L. (1987). Growth and development of the florunner peanut cultivar as influenced by plant population, planting date and water availability. *Peanut Sci*, 14, 11- 16.
18. Mixon, A. S. (1989). Influence of irrigation, row spacing on yield and market quality of peanut. *Applied Agricultural Research*, 1(5), 289- 293.
19. Mazing, R. W. & Coffelt, T. A. (1984). Row pattern and seedling rate effects on value of Virginia type peanut. *Agron J*, 76, 460- 462.
20. Munda, D. C. & Patel, C. S. (1989). Date of sowing, spacing and nutrient requirement of groundnut (*Arachis hypogaea* L.) under mid- altitude of Meghalaya. *Indian Journal of Agricultural Sci*, 59(11), 706- 709.

(In Farsi).

21. Wells, T., Anderson, W. F. & Wynne, J. C. (1991). Peanut yield as a result of fifty years of breeding. *Agron J*, 83, 937- 961.
22. Widaryanto, E. (1994). Influence of preemergence doses of the herbicide oxyfeloerfen (Goal 2 E) and population density of peanut (*Arachis hypogaea* L.) on weed control and its effect on growth and yield of peanut plants on dry land. In: *Proceedings of American Peanut Research and Education Society*. 12: 36.
23. Williams, J. H., Ndunguru, B. J. & Greenberg, D. C. (1995). Assessment of groundnut cultivars for end- of season drought tolerance in saheran environment. *Journal of Agricultural Science*, 125, 79- 85.
24. Wynne, J. C. (1974). *Breeding potential of the subspecies of Arachis hypogaea L. inheritance of branching pattern, estimates of combining ability and photoperiodic response*. Ph. D. Diss. North Carolina State Univ., Raleigh, NC. 45- 56.
25. Yilmaz, H. A. (1999). Effect of different plant densities of two groundnut (*Arachis hypogaea* L.) genotype on yield, yield components and oil and protein contents. *Turkish Journal of Agriculture and Forestry*, 23, 299- 308.