
(*Saccharomyces cerevisiae*)

(*Artemia urmiana*)

*

(/ / : / / :)

(*Saccharomyces cerevisiae*)

(*Artemia urmiana*)

() / () / () / () /
()

(/ ± /) (/ ± /) (/ ± /)
/ ± /) .(P< /) (/ ± /)
(/ ± /) (/ ± /) (/ ± /)
/ .(P< /)

(*Saccharomyces cerevisiae*)

(*Artemia urmiana*)

:

...

(Soltanian *et al.*, 2007 2011

mnn9

(Sorgeloos *et al.*, 1986)

-

(Marques *et al.*, 2005)

Marques *et al.*,)

(2005; Soltanian *et al.*, 2007

/

-

/

(Zmora *et al.*, 2002)

Dunaliella

Chaetoceros Tetraselmis sochrysis

Sajeevan)

Scenedesmus Spirulina

-

(*et al.*, 2006

-

(Verschuere *et al.*, 1997)

(Suphantharika *et al.*, 2003; Sajeevan *et al.*, 2009)

(Marques *et al.*, 2005)

-

BOD

Marques *et al.*,)

(2005

Ahmadnia *et al.*,)

Agh *et al.*,) / () /
 () / () / ()
 (2007) (S. cerevisiae)
 pH)
 (

() /
 () / () / () /

(Warland and Warland, 2001))
 / pH (

(Warland and Warland, 2001)

(Lavens and Sorgeloos, 1996)

(Verschuere *et al.*, 1997) pH

$$1000 \times [(\text{طول آرتمیا لگاریتم}) + (\text{طول آرتمیا لگاریتم}) \times 2.5] = \text{وزن خشک آرتمیا (میکروگرم)}$$

$$() = \frac{1}{100} \times \times$$

(Farhangi and Carter, 2001)


...

(ANOVA) $[(\quad - \quad) / \quad] =$
SPSS $P < /$ $\text{Ln} \quad) \times =$
Excel $[\quad / (\quad)] \times =$ (Ln



$\times \quad / \quad =$

Excel

Arcsin




(/) (/) (n=) () (±)
() / (/)



...

(n =)		(±)	
a / ± /	a / ± /	a / ± /	a / ± / ()
a / ± /	a / ± /	b / ± /	a / ± / ()
a / ± /	ab / ± /	c / ± /	b / ± / ()
a / ± /	a / ± /	b / ± /	a / ± /
a / ± /	a / ± /	b / ± /	a / ± / ()
a / ± /	b / ± /	c / ± /	b / ± /
c / ± /	ab / ± /	a / ± /	a / ± /

*



(Thinh *et al.*, 1999)

(/ /)
 (/ /)
 (/ /)
 S.
Americanuphis Nereis diversicolor
 (*Glycera* spp.) (*reesei*)
 (Naegel, 1999)
cerevisiae

.(Naegel, 1999)

$l \pm l$ *Chaetoceros*

.(Juarez Carrillo, 1995)

Chaetoceros

.(Naegel, 1999) $l \pm l$) ($l \pm l$)

) (

(

(Coutteau *et al.*,

Dunaliella tertiolecta 1990)

.(Vanhaecke and Sorgeloos, 1989) () B_2

$l \pm l$ (Yamada and Sgarbieri, 2005)

- Li) ()

and Gatlin III, 2003; Yamada and Sgarbieri, .(2005

(Mannoprotein)

(Coutteau *et al.*, 1990)

/

$l \pm l$)

(

.(Naegel, 1999) .(Peter *et al.*, 1990)

/

(*Artemia salina*) (

Dunaliella tertiolecta

Mason,)

.(1963 BOD

-

(Naegel, 1999)

(Marques *et al.*, 2005)

-

(Lashkarizadeh, 2010)

S.)

/ (Cerevisiae × Chaetoceros) /
 (Naegel, 1999) /
 / () /

References

- Agh, N., Abatzopoulos, T.J., Kappas, I., Van Stappen, G., Razavi Rouhani, S.M. and Sorgeloos, P., 2007. Coexistence of sexual and parthenogenetic Artemia populations in lake Urmia and neighbouring lagoons. International Review of Hydrobiologia 92, 48-60.

- Ahmadnia, H.R., Farhangi, M., Rafiee, GH., Noori, F., Safari, O., 2011. Effects of different levels of Bacillus subtilis and Bacillus licheniformis on digestive enzyme activities in Artemia urmiana. Asian-Pacific Aquaculture . Kochi, India, P14.

- Coutteau, P., Lavens, P. and Sorgeloos, P., 1990. Baker's yeast as a potential substitute for live algae in aquaculture diets: Artemia as a case study. Journal of World Aquaculture Society 21 (1), 1-8.

- Farhangi, M. and Carter, C.G., 2001. Growth, physiological and immunological responses of rainbow trout (Onchorhynchus mykiss) to different dietary inclusion levels of dehulled lupin (lupins angustifolius). Aquaculture Research 32, 329-340.

- Juarez Carrillo, E., 1995. Evaluacion nutricional de Artemia franciscana, Kellogg, 1906, (Crustacea: Branchipoda) alimentada con Chaetoceros sp. enriquecido con fosfolipidos en emulsion. MSc thesis. from the Center of Scientific Investigation and Superior Education of Ensenada (CICESE), Ensenada, B.C. Mexico. 144 pp. (In French)

- Lashkarizadeh, M., 2010. Estimation of the nutritional requirements of Artemia urmiana until adult stage using the body composition and determination of the best diet in the terms of biomass, nutritional value and

the investigation of digestive enzyme changes. Department of Fisheries, Faculty of Natural Resources, University of Tehran, Iran. 159 pp. (In Persian)

- Lavens, P. and Sorgeloos, P., 1996. Manual on the Production and Use of Live Food for Aquaculture, University of Ghent, Ghent, Belgium, 295 pp.

- Marques, A., Dinh, T., Ioakeimidis, C., Huys, G., Swings, J., Verstraete, W., Dhont, J., Sorgeloos, P. and Bossier, P., 2005. Effects of bacteria on *Artemia franciscana* cultured in different gnotobiotic environments. *Applied and Environmental Microbiology* 71, 4307-4317.

- Mason, D., 1963. The growth response of *Artemia salina* (L.) to various feeding regimes. *Crustaceana* 5, 138-150.

- Naegel, L.C.A. 1999. Controlled production of Artemia biomass using an inert commercial diet, compared with the microalgae *Chaetoceros*. *Aquacultural Engineering* 21, 49-59.

- Orozco-Medina, C., Maeda-Martinez, A.M. and Lopez-Cortes, A., 2002. Effect of aerobic Gram-positive heterotrophic bacteria associated with *Artemia franciscana* cysts on the survival and growth. *Aquaculture* 213, 15-29.

- Peter, C., Patrick, L. and Patrick, S., 1990. Baker's yeast as a potential substitute for live algae in aquaculture diets: Artemia as a case Study. *Journal of the World Aquaculture Society* 21, 1-9

- Sajeevan, T.P., Philip, R. and Singh, I.S. B., 2006. Immunostimulatory effect of a marine yeast *Candida sake S165* in *Fenneropenaeus indicus*. *Aquaculture* 257, 150-155.

- Sajeevan, T.P., Philip, R. and Bright Singh, I.S., 2009. Dose/frequency: A critical factor in the administration of glucan as immunostimulant to Indian white shrimp *Fenneropenaeus indicus*. *Aquaculture* 287, 248-252.

- Soltanian, S., Dhont, J., Sorgeloos, P. and Bossier, P., 2007. Influence of different yeast cell-wall mutants on performance and protection against pathogenic bacteria (*Vibrio campbellii*) in gnotobiotically-grown Artemia. *Fish and Shellfish Immunology* 23, 141-153.

- Sorgeloos, P., Lavens, P., Léger, P., Tackaert, W. and Versichele, D., 1986. Manual for the culture and use of brine shrimp artemia in aquaculture. State University of Ghent, Belgium. Belg, pp. 319

- Suphantharika, M., Khunrae, P., Thanardkit, P. and Verduyn, C., 2003. Preparation of spent brewer's yeast beta-glucans with a potential application as an immunostimulant for black tiger shrimp, *Penaeus monodon*. *Bioresource Technology* 88, 55-60

- Think, L.-V., Renaud, S.M. and Parry, D.L., 1999. Evaluation of recently isolated Australian tropical microalgae for the enrichment of the dietary value of brine shrimp, artemia nauplii. *Aquaculture* 170,161-173.

- Vanhaecke, P. and Sorgeloos, P., 1989. International study on *Artemia*. XLVII. The effect of temperature on cyst hatching, larval survival and biomass production for different geographical strains of brine shrimp *Artemia* spp. *Annals Social Research of Zoology* 118, (1), 7-23.

- Verschuere, J.D.P.S.W.V., 1997. Monitoring Biolog patterns and r/K-strategists in the intensive culture of Artemia juveniles. *Journal of Applied Microbiology* 83, 603-612.

- Warland, D. and Warland, T., 2001. Artemia: decapsulation, hatching, feeding, ongrowing and enrichment, Oz Reef, Online, viewed 20 August 2008, <http://www.ozreef.org>.

- Yamada, E.A. and Sgarbieri, V.C., 2005. Yeast (*Saccharomyces cerevisiae*) protein concentrate: preparation, chemical composition and nutritional and functional properties. *Journal of Agricultural and Food Chemistry* 53, 3931-3936.

- Zmora, O., Avital, E. and Gordin, H. 2002 Results of an attempt for mass production of Artemia in extensive ponds. *Aquaculture* 213, 395-400.

Effects of Different Levels of Yeast (*Saccharomyces cerevisiae*) Feeding on Growth Factors and Survival of *Artemia urmiana*

H. R. Ahmadnia*¹, M. Farhangi¹, Gh. Rafiee¹ and P. Fazli²

¹ Department of Fisheries, Faculty of Natural Resources, University of Tehran, Karaj, I.R. Iran

² Faculty of Marine Science and Technology, Islamic Azad University, North Branch, Tehran, I.R. Iran

(Received: 24/12/2010 , Accepted: 18/10/2011)

Abstract

Artemia plays a major role in aquaculture as one of the most important live foods used in larval rearing. The aim of this study was to determine the optimum level of feeding in *Artemia urmiana* with Baker's yeast (*Saccharomyces cerevisiae*). Triplicate groups were fed with different levels of *S. cerevisiae* yeast 0.625 (control), 1.25 (T₁), 2.50 (T₂), 5.0 (T₃) (mg yeast per 1000 nauplii) for 10 days. Culture conditions such as temperature (29°C), Dissolved oxygen (5 mg/l), Salinity (60 g/l) and pH (8) were set according to Standard Method. At the end of the trial, total length, dry weight, survival rate and biomass production were measured. Moreover, average daily growth (ADG), body weight increase (BWI), specific growth rate (SGR) and condition factor (CF) were determined. The results showed that total length (2.10±0.14 mm), dry weight (11.75±1.91µg), biomass production (129.55±21.3 mg/l) and survival rate (62.2±12.2 %) of T₂ were significantly higher than. Also ADG (1.12±0.19 mg), BWI (11.21±1.89) and SGR (30.71±1.41) were significantly higher in T₂ compared to other treatments. The highest CF was observed in T₄ (60.47±21.46) and the lowest was observed in T₂ (22.89±3.01) (P<0.05). According to these results, the best level of *Artemia* feeding with *S. cerevisiae* was 1.25 mg of *S. cerevisiae* per 1000 *Artemia nauplii*. Also our results indicated that food efficiency depends on the amount of food available, and it was increased with food availability to a certain level.

Keywords: *Artemia urmiana*, Optimum feeding, Baker yeast (*Saccharomyces cerevisiae*), Growth factors, Survival