

***In vitro* antifungal activity of *Zataria multiflora* essential oil, fluconazole and ciclopirox olamine against *non-albicans Candida* species isolated from recurrent vulvovaginal candidiasis**

Mohammad Ali Esfandiary¹, Shahram Mahmoudi¹, Farideh Zaini¹, Mahin Safara¹, Mahmoud Mahmoudi², Alireza Farasat³, Azar Sabokbar⁴, Sanaz Abbasi⁴, Mohammad Ghahri⁵, Mohammad Amin Esfandiary⁶, Parivash Kordbacheh^{1*}

¹Department of Medical Parasitology and Mycology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran.

²Department of Epidemiology and Biostatistics, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran.

³Young Researchers & Elite Club, Pharmaceutical Sciences Branch, Islamic Azad University, Tehran, Iran

⁴Department of Microbiology, Karaj Branch, Islamic Azad University, Karaj, Iran.

⁵Department of Biological Sciences, College of Sciences, Imam Hussein University, Tehran, Iran.

⁶Faculty of pharmacy, Tehran University of Medical Sciences, Tehran, Iran.

* Corresponding Author: Email: pkordbacheh@tums.ac.ir , Tel: +98 21 88951583, Fax: +98 21 88951392

(Received: 25 March 2015, Accepted: 15 April 2015)

Abstract:

The aim of the present study was to investigate the antifungal activity of *Zataria multiflora* essential oil, fluconazole and ciclopirox olamine against 44 *non-albicans Candida* isolates from patients with recurrent vulvovaginal candidiasis. A set of *C. glabrata* (29 strains), *C. krusei* (3 strains) and *C. parapsilosis* (2 strains) were studied using Clinical and Laboratory Standards Institute (CLSI) broth macrodilution method. The results revealed that 33 isolates were resistant (MIC=64 µg/ml), 4 isolates were susceptible (MIC≤ 8 µg/ml) and 7 isolates had dose-dependent susceptibility (MIC= 16, 32 µg/ml) to fluconazole, respectively. With regard to fluconazole, high resistance rate was observed in *C. glabrata* and *C. krusei*. However, ciclopirox olamine was found to inhibit the growth of all *non-albicans Candida* species (MIC≤8 µg/ml). In this study, favorable antifungal activity against *non-albicans Candida* species was obtained by *Z. multiflora* despite having a wide range of MICs (34875-139500 µg/ml). The results indicated that ciclopirox olamine and *Z. multiflora* might be promising in the treatment of recurrent vulvovaginal candidiasis.

Keywords: ciclopirox olamine, fluconazole, *non-albicans candida* spp., recurrent vulvovaginal candidiasis, *Zataria multiflora*.

Introduction

Candida species are the etiologic agents of 20 to 25% of vulvovaginal infections.

Vulvovaginal candidiasis (VVC) is characterized by purities, burning, vaginal discharge, dyspareunia and dysuria. This

condition is uncommon before puberty, and at 25 years of age, approximately 50% of females must have had at least one episode of VVC and 75% of women must have experienced at least one episode of VVC during their lifetime (Trama *et al.*, 2005). Recurrent vulvovaginal candidiasis (RVVC) is defined as four or more episodes of *Candida* vaginal infection during a year and is estimated to occur in 5 to 8% of women during their fertility period (Sobel, 2006). Various epidemiologic surveys revealed that *C.albicans* is the most common etiologic agent of VVC, accounting for up to 85% of cases. However, some reports suggest that non-*albicans Candida* (NAC) species are responsible for 10 to 30% of episodes in specified geographical areas (Kennedy and Sobel, 2010). The evidence is slowly being aggregated on the increasing incidence of VVC owing to NAC species, especially *C.glabrata*, *C.tropicalis* and *C. krusei* (Faro, 1996).

Ciclopirox olamine (CPO) is a synthetic antifungal agent. It is a hydroxypyridone derivate and is different in mechanism of action and structure from the other recognized antifungal drugs. Its action may be via the chelation of trivalent metal cations (Fe^{3+}). CPO also prevents metal-dependent enzymes, such as peroxidase and catalase, which play an important role in the intracellular destruction of poisonous peroxides. Furthermore, CPO shows some antibacterial activities (Leem *et al.*, 2003). *Z. multiflora* (family: Lamiaceae) is a plant that usually grows in Iran, Afghanistan and Pakistan. The major components of the essential oil of this plant are phenolic compositions such as carvacrol and thymol (Khosravi *et al.*, 2009).

Since women with RVVC present a difficult management problem and these patients often suffer from depression and psychosexual problems, therefore specific attention to diagnosis and management of these patients are important. This study was conducted to evaluate the anti *Candida* activities of *Z.*

multiflora essential oil (ZMEO) and CPO against NAC species isolated from patients with RVVC in order to find an effective drug for the treatment of RVVC.

Materials and Methods

Fungal isolates: A total of 44 NAC species (*C.glabrata*; 29, *C.kefyer*; 10, *C.krusei*; 3 and *C. parapsilosis*; 2) from patients with RVVC were included in this study. *C. parapsilosis* ATCC22019 (fluconazole MIC ranges: 2–8 μ g/ml) and *C.krusei* ATCC6258 (fluconazole MIC ranges: 16–64 μ g/ml) were used as standard control species.

***In vitro* antifungal susceptibility testing:** Antifungal susceptibility tests were performed using the Clinical Laboratory Standards Institute (CLSI) M27-A2 broth macrodilution method (CLSI, 2008). RPMI 1640 (Gibco, USA) culture medium buffered at pH=7.0 with MOPS (Sigma-Aldrich, USA) was used for broth susceptibility testing. Water and 100% dimethyl sulfoxide (Merck, Germany) were used to dissolve fluconazole (Sigma-Aldrich, USA), ciclopirox olamine (USP, USA) and ZMEO (Barijessence, Iran), respectively. Two-fold serial dilutions of Flu, CPO and ZMEO were prepared in RPMI 1640 to achieve a concentration of 0.03125-64 μ g/ml for Flu, 0.00195-8 μ g/ml for CPO and 1089.8-558000 μ g/ml for ZMEO. For yeast inoculums preparation, two to three colonies (diameter >1 mm) of an overnight culture of *Candida* spp. on Sabouraud agar (Merck, Germany) at 35-37°C were suspended in 2 ml normal saline. The cell density was adjusted to $1-5 \times 10^4$ CFU/ml using hemocytometer slide. According to M27-A2 standard procedure of CLSI, each macroplate had two drug free controls, one with the media alone and the other with the media containing an equivalent amount of suitable solvent used to dissolve the drug. Plates were incubated at 35-37°C and results were read visually after 48 h and susceptibility levels were defined according to the MIC breakpoints suggested by CLSI. All

tests were carried out at least twice.

Results

Based on the results of this study, it was demonstrated that 28 out of 29 isolates of *C.glabrata* (MIC=64 µg/ml), 3 out of 10 isolates of *C. kefyer* (MIC=64 µg/ml) and 2 out of 3 isolates of *C. krusei* (MIC=64 µg/ml) were resistant to Flu. Susceptibility to Flu was seen in a few number of isolates including 1 isolate of *C.glabrata* (MIC=4 µg/ml), 2 isolates of *C. kefyer* (MIC= 4 and 8 µg/ml) and 1 isolate of *C. parapsilosis* (MIC=8 µg/ml). From all 44 NAC isolates, dose dependent susceptibility to Flu was seen in 5 isolates of *C. kefyer* (MIC=16-32 µg/ml), 1 isolate of *C.krusei* (MIC=16 µg/ml) and 1 isolate of *C.parapsilosis* (MIC=16 µg/ml) (Additional information is shown in Table 1).

In addition, in almost all 44 NAC isolates, MIC of CPO was clearly lower than Flu. However, MIC was 8 µg/ml for 42 out of the 44 isolates that includes *C.glabrata* (29 isolates), *C. kefyer* (8 isolates), *C.krusei* (3 isolates) and *C.parapsilosis* (2 isolates). For the 2 remaining *C. kefyer* isolates, the MIC of CPO was 4 µg/ml.

The growth of 30 isolates, which includes *C. glabrata* (17 isolates), *C. kefyer* (8 isolates), *C. krusei* (3 isolates) and *C. parapsilosis* (2 isolates), were inhibited at 69750 µg/ml of ZMEO. In the case of four (4) *C. glabrata* and one (1) *C. kefyer* isolates, the MIC of ZMEO was 34875µg/ml, and for the other remaining eight (8) *C. glabrata* and one (1) *C. kefyer* isolates, MIC was 139500 µg/ml.

Table 1. GM, MIC 50, MIC 90 and MIC ranges of fluconazole, Ciclopirox olamine and Zataria multiflora essential oil against C.glabrata, C.kefyer, C.krusei and C.parapsilosis isolates from recurrent vulvovaginal candidiasis

<i>Candida species</i> (number of isolates)	Antifungal concentration (µg/ml)			
	Flu	CPO	ZM	
	GM	58.16468	8	76747.7391
<i>C. glabrata</i> (29)	MIC 50	64	8	69750
	MIC 90	64	8	139500
	MIC range	4-64	8	34875-139500
	GM	21.1121	6.9644	69750
<i>C. kefyer</i> (10)	MIC 50	16	8	69750
	MIC 90	64	8	69750
	MIC range	4-64	4-8	34875-139500
	GM	40.3175	8	69750
<i>C. krusei</i> (3)	MIC 50	64	8	69750
	MIC 90	64	8	69750
	MIC range	16-64	8	69750
	GM	11.3137	8	69750
<i>C. parapsilosis</i> (2)	MIC 50	8	8	69750
	MIC 90	16	8	69750
	MIC range	8-16	8	69750

Flu: Fluconazole - CPO: Ciclopirox olamine – ZM: Zataria multiflora – GM: Geometric Mean - MIC 50, MIC 90: Minimum Inhibitory Concentration which inhibits 50 and 90 % of isolates, respectively.

Discussion

Treatment of RVVC still faces difficulties and is yet to be determined. The incidence of RVVC due to NAC species is increasing significantly in recent years. These species have also shown significant drug resistance (Sobel, 2006; Watson and Pirotta, 2011).

The results of the present study revealed a high resistance to Flu among NAC species isolated from RVVC. On the other hand, high antifungal activity of CPO presented its potency as a good alternative choice for the treatment of patients with Flu resistant NAC species infections. Also growth inhibition of 30 out of 44 NAC isolates by ZMEO was good evidence to prove the notable antifungal activity of this medicinal plant.

Furthermore, the result of the present study revealed high resistance to Flu among *C. glabrata* isolates (28 out of 29) with 4-64 µg/ml MIC range. This outcome is inconsistent with the results of Padua, Asticcioli and Pfaller with MIC ranges of 0.25-4, 0.125-64 and 0.5-128 µg/ml, respectively (Pádua *et al.*, 2003; Asticcioli *et al.*, 2009; Pfaller *et al.*, 2011). However, the results of Richter (MIC range=2-64 µg/ml), and Badiee (MIC range=8-64 µg/ml) are almost similar to the findings of this study (Richter *et al.*, 2005; Badiee *et al.*, 2010).

However, as shown in this study, the MIC range of Flu against *C. Kefyer* isolates (4-64 µg/ml) was considerably higher than those achieved by Badiee and Ozcelik (0.25-1 µg/ml and ≤0.03-4 µg/ml, respectively) (Badiee *et al.*, 2010; Ozçelik *et al.*, 2006).

In addition, the susceptibility of *C. krusei* and *C. parapsilosis* to Flu in this study's results were not similar to the studies conducted by Richter, Ozcelik, Satana and Swinne (Richter *et al.*, 2005; Ozçelik *et al.*, 2006; Satana *et al.*, 2010; Swinne *et al.*, 2005).

CPO is a synthetic drug with broad-spectrum anti-fungal properties (Gupta and Plott, 2004). The results of this study revealed lower MIC ranges of CPO against all previous studied NAC-species in comparison to Flu. This could be a therapeutic efficacy indicator of CPO in the treatment of RVVC owing to the NAC-species. In spite of this, the values of MICs for CPO in the present study were considerably higher than those reported by Hanel, Czaika and Gupta (Hanel *et al.*, 1988; Czaika *et al.*, 2000; Gupta and Kohli, 2003). Broad spectrum antimicrobial activity, high mucosal absorption, multiple mechanisms of antifungal activity and fewer side effects are some of CPO advantages in the treatment of vaginal infections (Zaneveld *et al.*).

The results of the present study revealed generally higher MIC ranges for Flu and CPO against NAC-species (*C. glabrata*, *C. kefyer*, *C.krusei* and *C. parapsilosis*). This may be attributed to the higher drug resistance of investigated isolates leading to recurrent vulvovaginal symptoms in patients with RVVC.

In traditional medicine, *Z. multiflora* has been used as a medicinal plant in the treatment of dysmenorrhea, spasms, respiratory infections and digestive disorders (Simbar *et al.*, 2008). Also, anti inflammatory and analgesic effects of this plant have been reported (Hosseinzadeh *et al.*, 2003). Owing to limited geographical distribution of *Z. multiflora*, there are few studies on its *in vitro* antifungal activity against NAC-species. Mahmoudabadi evaluated the *in vitro* anti *Candida* activity of aqueous, ethanolic and methanolic extract of *Z. multiflora* against *albicans* and non- *albicans Candida* species. Their results showed a notable anti *Candida* activity for methanolic followed by ethanolic extract. Aqueous extract has no anti *Candida* activity, and *C. parapsilosis* was the most

sensitive species (Mahmoudabadi *et al.*, 2006). Moreover, Zomorodian showed a considerable anti *Candida* activity for *Z. multiflora* with MIC value ranging from 0.07 to 0.5 µl/ml (Zomorodian *et al.*, 2011). Gandomi reported that *Z. multiflora* mainly affects the cell membrane and cell wall of fungi. This may be a mechanism for the antifungal activity of *Z. multiflora* (Gandomi *et al.*, 2011). Khosravi studied the efficacy of 0.1% *Z. multiflora* cream in the treatment of acute vaginal candidiasis in comparison to 1% clotrimazole. The results showed 90 and 74.8% recovery in patients that received *Z. multiflora* cream and clotrimazole, respectively (Khosravi *et al.*, 2008). The results of the present and previous studies reveals the powerful potency of *Z. multiflora* as a natural antifungal agent. Therefore, more studies need to be conducted in order to identify the effective compound of this medicinal plant.

Conclusion

The results of numerous studies conducted across the globe on the drug susceptibility of NAC species, revealed different patterns of drug resistance, in various geographical areas. Therefore, more studies should be conducted in different parts of the world. The favorable activity of CPO and ZMEO against NAC isolates in this study implies their efficacy in the treatment of RVVC in Iranian patients.

Acknowledgments

This research was supported by grant NO. 240.3513 89.10.5 of Tehran University of Medical Sciences.

The authors are grateful to the Razak Laboratories Co. staff, especially Dr. Karbassi, Dr. Jabbarian, Mrs. Jannati, Mr. Arian, Mr. Bakshi and Barijessence Company for their co-operation.

References

1. Asticcioli, S., Sacco, L., Daturi, R., Matti, C., Nucleo, E., Zara, F., Pagani, L., 2009. Trends in frequency and in vitro antifungal susceptibility patterns of *Candida* isolates from women attending the STD outpatients' clinic of a tertiary care hospital in Northern Italy during the years 2002-2007. *New Microbiol*, 32(2): 199-204.
2. Badiie, P., Alborzi, A., Davarpanah, M.A., Shakiba, E., 2010. Distributions and Antifungal Susceptibility of *Candida* Species from Mucosal Sites in HIV Positive Patients. *Archives of Iranian medicine*, 13(4): 282-287.
3. Clinical and Laboratory Standards Institute. Reference method for broth dilution antifungal susceptibility testing of yeasts; Approved standard, 3rd ed, CLSI document M27-A3, Clinical and Laboratory Standards Institute, Wayne, PA. USA. (2008).
4. Czaika, V., Tietz, H.J., Schmalreck, A., Sterry, W., Schultze, W., 2000. Antifungal susceptibility testing in chronically recurrent vaginal candidosis as basis for effective therapy. *Mycoses*, 43: 45-50.
5. Faro S., 1996. New treatments for vulvovaginal candidiasis. *Infect dis obstet gynecol*, 4(4): 247-254.
6. Gupta, A.K., Plott, T., 2004. Ciclopirox: a broad-spectrum antifungal with antibacterial and anti-inflammatory properties. *Int J Dermatol*, 43 (1): 3-8.
7. Gupta, A.K., Kohli, Y., 2003. In vitro susceptibility testing of ciclopirox, terbinafine, ketoconazole and itraconazole against dermatophytes and nondermatophytes, and in vitro evaluation of combination antifungal activity. *Br J Dermatol*, 149(2): 296-305.
8. Hanel, H., Raether, W., Dittmar, W., 1988. Evaluation of fungicidal action in vitro and in a skin model considering the influence of penetration kinetics of various standard antimycotics. *Ann NY Acad Sci*, 544: 329-337.
9. Hosseinzadeh, H., Ramezani, M., salmani, G., 2000. Antinociceptive, anti-

- inflammatory and acute toxicity effects of Zataria multiflora Boiss. extract in mice and rats. *J Ethnopharmacol*, 73: 379-385.
10. Gandomi, H., Misaghi, A., Akhondzadeh Basti, A., Hamed, H., Ramezani Shirvani, Z., 2011. Effect of Zataria multiflora Boiss. essential oil on colony morphology and ultrastructure of *Aspergillus flavus*. *Mycoses*, 54 (5) : 429-437.
 11. Kennedy, M.A., Sobel, J.D., 2010. Vulvovaginal candidiasis caused by non-albicans Candida species: new insights. *Curr Infect Dis Rep*, 12(6): 465-470.
 12. Khosravi, A.R., Shokri, H., Tootian, Z., Alizadeh, M., Yahyaraeyat, R., 2009. Comparative efficacies of Zataria multiflora essential oil and itraconazole against disseminated Candida albicans infection in BALB/c mice. *J.Microbiol*, 40(3): 439-445.
 13. Khosravi, A.R., Eslami, A.R., Shokri, H., Kashanian, M., 2008. Zataria multiflora cream for the treatment of acute vaginal candidiasis. *Int J Gynaecol Obstet*, 101(2) : 201-2.
 14. Leem, S.H., Park, J.E., Kim, I.S., Chae, J.Y., Sugino, A., Sunwoo, Y., 2003. The possible mechanism of action of ciclopirox olamine in the yeast *Saccharomyces cerevisiae*. *Mol Cells*. 15:55-61.
 15. Mahmoudabadi, A.Z., Dabbagh, M.A., Fouladi, Z., 2006. In vitro anti-Candida activity of Zataria multiflora boiss. *eCAM*, 4: 351-353.
 16. Ozçelik, B., Balaban, N., Aksaray, S., Cesur, S., Kaynak, F., Çayırılı, A., 2006. In-vitro Susceptibility of *Candida* spp. Isolated from Clinical Specimens Against some Antifungal Agents. *Turkish J Pharm Sci*, 3(1): 8-16.
 17. Pádua, R.A.F., Guilhermetti, E., Svidzinski, T.I.E., 2003. In vitro activity of antifungal agents on yeasts isolated from vaginal secretion. *Acta Scientiarum*, 25: 51-54.
 18. Pfaller, M.A., Hata, K., Jones, R.N., Messer, S.A., Moet, G.J., Castanheira, M., 2011. In vitro activity of a novel broad-spectrum antifungal, E1210, tested against *Candida* spp. as determined by CLSI broth microdilution method. *Diagn Microbiol Infect Dis*, 71(2): 167-170.
 19. Richter, S.S., Galask, R.P., Messer, S.A., Hollis, R.J., Diekema, D.J., Pfaller, M.A., 2005. Antifungal susceptibilities of *Candida* species causing vulvovaginitis and epidemiology of recurrent cases. *J Clin Microbiol*, 43(5): 2155-2162.
 20. Satana, D., Genc, G.E., Erturan, Z., Afr, J., 2010. The antifungal susceptibilities of oral *Candida* spp. isolates from HIV-infected patients. *J Microbiol Res*, 4(17): 1831-1835.
 21. Simbar, M., Azarbad, Z., Mojab, F., Alavi Majd, H.A., 2008. Comparative Study of Therapeutic Effects of Zataria Multiflora Vaginal Cream and Metronidazole Vaginal Gel on Bacterial Vaginosis. *Phytomedicine*, 13(3): 193-202.
 22. Sobel, J.D., 2006. Management of recurrent vulvovaginal candidiasis. *Curr Infect Dis Rep*, 8(6): 481-486.
 23. Swinne, D., Watelle, M., Nolard, N., 2005. In vitro activities of voriconazole, fluconazole, itraconazole and amphotericin B against non *Candida albicans* yeast isolates. *Rev Iberoam Micol*. 22(1): 24-28.
 24. Trama, J.P., Adelson, M., Raphaelli, I., Stemmer, S., Mordechai, E., 2005. Detection of *Candida* species in vaginal samples in a clinical laboratory setting infect. *Dis.Obstet Gynecol*, 13: 63-67.
 25. Watson, C., Pirotta, M., 2011, Recurrent vulvovaginal candidiasis current management. *Aust Fam Physician*, 40(3): 149-151.
 26. Zaneveld, L.J.D., Waller, D.P., Sellors, J., Camus-Bablon, F., Ciclopirox olamine: A vaginal product with Microbicide potential. Available on www.path.org.
 27. Zomorodian, K., Saharkhiz, M.J., Rahimi, M.J., Bandegi, A., Shekarkhar, G., Bandegani, A., Pakshir, K., Bazargani, A., 2011. Chemical composition and antimicrobial activities of the essential oils from three ecotypes of Zataria multiflora. *Pharmacognosy magazine*, 7(25): 53-59.