

A Review of on Environmental Pollution Bioindicators

Asif, N^{*}., Malik, M.F. and Chaudhry, F.N.

Department of Zoology, University of Gujrat, Hafiz Hayat Campus, Gujrat,
Pakistan

Received: 09.07.2017

Accepted: 04.10.2017

ABSTRACT: Qualitative status of the environment is signaled by a group of indicators, known as bioindicators, several of which are responsible for showing progressive impacts of different types of pollutants. Having addressed the influence of various bioindicators in environmental pollution, it has been revealed that bioindicators are sensitive to any disturbance in any environment. With regards to the pollution, the quality of an ecosystem can be judged by an organism, which is actually an indicator and play a key role in monitoring its changes. A reliable and cost effective way to evaluate the changes in the environment is possible by means of indicator species as ecological indicators, yet selecting a specific indicator poses a real challenge, followed by its identification as well as relation among indicators and their particular applications. As a result, environmental, ecological, and biodiversity indicators fulfill their goal of monitoring environmental quality. The current situation requires cost effective bioindicators along with their reliability to detect and mitigate the impacts of pollution in our environment.

Keywords: Bioindicators, Indicator Species, Environmental Pollution

INTRODUCTION

Bioindicator is given to a living entity or group of organisms that shows the information, either based on the environment or a constituent of it (Wilkomirski, 2013). Keeping this definition in mind, the current study aims to select different types of bioindicators such as microorganisms, lichens, animals, or plants which under environmental alterations tend to produce certain molecular signals (Posudin, 2014). There have been New fields of research due to the invasion of a wide range of individual components in the environment in toxicological, chemical, and ecological terms (Merian et al., 2008).

Complete monitoring of the whole area

is possible by bioindication, which indicates various living systems with simple data (Mueller, 1980). The effect of external factors on ecosystems can be assessed by reliable procedure of bioindication (Markert, 2008). Living organisms that can be examined without any difficulty and environmental conditions of their habitat can be regarded as indicator species (Landres et al., 1988; Cairns and Pratt, 1993; Bartell, 2006; Burger, 2006). Environment renders indicator species sensitive to its alterations, whereas detection of ecosystem by assessing an efficient incentive of a single population is believed to be more useful and cheaper (Spellerberg, 2005). Short-term stress conditions or long-term events predict future situations and alterations by

*Corresponding author, Email: asifnayyab@gmail.com

identifying the variations in indicator species (Cairns & Pratt, 1993).

Several restrictions on indicator species have been described, despite their increased popularity (Lindenmayer et al., 2000; Lindenmayer & Fischer, 2003; Morrison, 2009; Lindenmayer & Likens, 2011). Environment complexity with single population rarely occurs and have subjective selection criteria for indicators according to initial restrictions (Ahmed et al., 2016). In addition, differing environmental indicators respond to various environmental alterations, e.g. ecological indicators demonstrate the changes in living systems' environment, whereas diversity coexists with the overall community diversity ecosystem for taxonomic groups of biodiversity indicators (Mc Geoch, 1998). Therefore, monitoring goals are separated by three categories, with regards to the changes in either physical or chemical changes of the environment along with biodiversity and ecological procedures (Holt & Miller, 2011). The present study shall discuss the impact of various bioindicators in environmental pollution.

Bat as Bioindicator of Environment Health Assessment

Human population is increasing at a frightening rate. Currently, population of over 7 billion humans helped as well as deteriorative effects against the balance of living entities and humans is devastating the earth,(Barnosky et al., 2012). To attain a balanced living environment, bioindicators such as bat play a vital role to lessen human impact for monitoring environmental health (McGeoch, 1998).

Among most diverse vertebrate groups, bat is one with more than 1300 species, sensitive to habitat deterioration and land use (Fenton & Simmons, 2014). Bats are cost effective, responsive to environment stressors, and stable taxonomically, providing a wide range of services from pollination to pest control in the ecosystem (Jones et al., 2009; Jones, 2012).

A key character of appropriate bioindicators like bat is to respond to alterations in an ecosystem, e.g. the ones in drought events (Amorim et al., 2015), agricultural practices (Park, 2015), urbanization (Ancillotto & Russo, 2015), light pollution (Stone et al., 2015), and heavy metals (Zukal et al., 2015).

Tourist Disturbance, Indicated by Birds and Fish

The basic factors to affect biodiversity of freshwater environments include over exploitation and pollution (Cooperrider & Noss, 1994; Curtis et al., 1998) with tourism being an important source of chaos in these environments (Palacio et al., 2007). Similar responses cannot be delivered by various indicators (Duelli & Obrist, 2003).

Although most common characters are shared by bioindicators such as fish (Pyrovetsi & Papastergiadou, 1992; Browder et al., 2002) and birds (Heino et al., 2005; Fu et al., 2003), like being short lived species after disturbance, some tourist activities may affect these groups as well as bioindicators of relative human disturbance (Tershy et al., 1997; Higginbottom et al., 2003; Newsome et al., 2004).

Freshwater Mussels as Biological Indicators

Changes in water and habitat have been recorded from the use of feasible indicator acts as environmental logbooks which are the properties of freshwater mussels (Ortmann 1909; Wurtz 1956; Bedford et al., 1968; Simmons & Reed, 1973; Imlay, 1982; Neves, 1993; Naimo, 1995).

Alterations in habitat are promoted by humans, an ordered damming of creeks and rivers has had the most significant effect on freshwater mussels (Bogan, 1993; Neves, 1993; Yeager, 1993). The physical, chemical, and biological attributes of numerous rivers have changed from shallow flowing habitats to long linear pools drastically (Ellis, 1942; Bates, 1962; Coon et

al., 1977; Yeager, 1993; Hughes & Parmalee, 1999).

Sedimentation is another process with harmful impacts on freshwater mussel communities. Soft, cohesive substrates, and suspended fine sediments are deleterious for most species and may affect respiration, feeding, and growth (Marking & Bills, 1979).

Honey Bee as Bioindicator of Environment Quality

Likely to be used to determine environmental quality for bioindication, honey bee is an efficient bioindicator that reacts quickly to various external factors (Crane, 1984; Bilalov et al., 1992; Jeliaskova et al., 2001; Jeliaskova et al., 2002; Porrini et al., 2003; Zhelyazkova et al., 2004; Fakhimzadeh et al., 2005; Stanimirovic et al., 2005; Bianu & Nica, 2006; Gallina et al., 2006; Spodniewska & Romaniuk, 2006).

Cases of environmental pollution and atmospheric air which have contributed imbalanced health level, life status, and quality of the population in the past few years, have been monitored (Berberova et al., 2008; Takuchev, 2011). The existing problem in the environment is monitored by determining the traces in plant and animal origins along with honey bees and humans (Eneva & Todorova, 2004; Berberova et al., 2008; Petkov et al., 2010).

As a consequence of atmospheric nuclear testing, bee has been monitored as an indicator of radionuclide strontium 90 in the environment (Svoboda, 1962). Since 1970, in territorial and urban surveys, environmental pollution is caused by heavy metals (Cavalchi & Fornaciari, 1983; Crane, 1984) and pesticides in rural regions (Atkins et al., 1981).

Earthworm as Bioindicator of Soil Pollution

Both environment and human life are in danger, due to high levels of pesticides and heavy metals, which cause soil pollution,

(Caroline et al., 2001) with the exposed organisms being the greatest invaders of such kind of complex effects (Svendsen et al., 2004). Among soil species, in most soils, a huge part of invertebrate biomass is formed by earthworms (Haeba et al., 2013).

All biological agent, crusher, moisture retainer, aerator, and nature's plough are both composting agents and biofertilizers, at the same time (Eguchi et al., 1995). Predators play a key role in the assimilation of contaminated earthworm tissues increase the level of harmful chemicals in food chain and in this way soil health is indicated by these biological indicators with their particular behavior in toxic soils (Caroline et al., 2001). By means of 'earthworm acute toxicity test' possible risk of environmental pollutants on invertebrates of soil has been examined (Anonymous, 1984).

Lichen as Bioindicator of Metal Pollution

Pb, Ni, Cu, Cr, and Cd are some metals in cement dust, generated by cement industry (Alkhasman & Shawabkeh, 2006). Wind and rain cause spread cement dust and its related chemicals usually found in lichens, soils, animals, and plants (Schuhmacher et al., 2009). To detect air pollution, lichens are likely to be used as indicators (Yazici & Aslan, 2006; Cicek et al., 2008; Bingol et al., 2009). Due to their sensitivity to different factors of environment, lichens are considered to be the most appropriate biomonitors of air quality during last 30 years (Conti & Cecchetti, 2001; Brodo, 1961; Rossbach & Lambrecht, 2006).

Lichens are considered the most reliable biomonitors according to their specific physiological, morphological, and anatomical characteristics (Battal et al., 2004). Changes in air quality can be detected by air pollutant sensitive epiphytic lichen species (Showman, 1988). It is compulsory for the judgment of an area that it should have enough species of lichen to monitor air pollution (Calvelo & Baccala, 2009).

Phytoplankton as Bioindicator for Water Quality

There is a relation between aquatic organisms and pollution as aquatic organisms are considered to be the indices of pollution (Kolkwitz and Marrson, 1908). Several methods have been proposed to locate organisms, able to monitor the quality of water (Knopp, 1954; Zelinka & Marvan, 1961; Sladeczek, 1973). In some streams of Taiwan, microorganisms, fish, and macroinvertebrates have been used to monitor water quality (Hau et al., 1976; Hong, 1979; Lee et al., 1967).

Both anthropogenic activities and industrialization are behind the hazardous materials and the pollutants, increasingly discharged in the environment (Ghorbanli et al., 2007; Raabe, 1999; Bakand et al., 2005; Hayes et al., 2007). Biomonitoring capacity of roadside plant leaves can be measured by their exposure to air pollutants as well as their reaction as stressor against them (Pandey et al., 2005; Sharma et al., 2007), yet in an industrial area the response from several growing plants has been monitored biochemically and physiologically through proper investigation (Joshi et al., 2009; Gupta et al., 2009; Sharma and Tripathi, 2009; Gupta et al., 2011).

CONCLUSION AND RECOMMENDATIONS

The use of indicator species as ecological indicators is reliable and cost-effective to assess the changes in environment, though it is quite challenging to select a specific indicator and then identify it as well as the relation among the indicators and their particular applications. The future usefulness of indicator species will strictly depend on selected indicator groups that present the environment in actually, showing efficient relations between the indicator species and under lying processes of interest.

Furthermore, extensive surveys, search of literature, and research is required on various bioindicators as well as their

particular role in determination of environmental health including novel approaches to human interference that creates disturbance in the environment, causing pollution and leading to the loss of ecosystem services such as plant pollinators, drinking water, and clean air.

Environmental pollution has major impacts on the disturbance of ecosystem. Although it is very difficult to make our environment free from pollution, it can be reduced by releasing chemical liquids from factories into water bodies after treatment, using vehicles with less fuel combustion, and using pesticide spray in a controlled manner. Moreover, further studies are required in this aspect to prevent living beings from suffering damages that play vital role in the maintenance of ecosystem.

REFERENCES

- Ahmed, A.H.S., Aaron, M.E., Alison, O., Claudia, V.L. and Matthew, K.L. (2016). How do ecologists select and use indicator species to monitor ecological change? Insights from 14 years of publication in *Ecological Indicators*. *Ecological Indicators*, 60, 223-230.
- Alkhashman, O.A. and Shawabkeh, R.A. (2006). Metals distribution in soils around the cement factory in southern Jordan. *Environ. Pollut.*, 140, 387-394.
- Amorim, F., Mata, V.A., Beja, P. and Rebelo, H. (2015). Effects of a drought episode on the reproductive success of European free-tailed bats, *Tadarida teniotis*, *Mammalian Biology*.
- Ancillotto, L. and Russo, D. (2015). Sensitivity of bats to urbanization: a review. *Mammalian Biology*, this issue.
- Anonymous, (1984). Guidelines for the testing of chemicals No. 207. Earthworm acute toxicity tests. OECD Adopted 4 April 1984.
- Atkins, E. L., Kellum, D. and Atkins, K.W. (1981). Reducing pesticides hazard to honey bees: mortality prediction techniques and integrated management strategies. Division of Agricultural Sciences, University of California, Leaf. 2883, Pp: 22.
- Bakand, S., Winder, C., Khalil, C. and Hayes, A. (2005). Toxicity assessment of industrial chemicals and airborne contaminants: Transition from in vivo to in vitro test method: A review. *Inhal.Toxicol.*, 17(13), 775-787.

- Barnosky, A.D., Hadly, E.A., Bascompte, J., Berlow, E.L., Brown, J.H., Fortelius, M., Getz, W.M., Harte, J., Hastings, A., Marquet, P.A., Martinez, N.D., Mooers, A., Roopnarine, P., Vermeij, G., Williams, J.W., Gillespie, R., Kitzes, J., Marshall, C., Matzke, N., Mindell, D.P., Revilla, E. and Smith, A.B. (2012). Approaching a state shift in Earth's biosphere. *Nature*, 486, 52-58.
- Bartell, S.M. (2006). Biomarkers, bioindicators, and ecological risk assessment, a brief review and evaluation. *Environ. Bioindic.*, 1, 39-52.
- Bates, J.M. (1962). The impact of impoundment on the mussel fauna of Kentucky Reservoir, Tennessee River. *American Midland Naturalist*, 68, 232-236.
- Battal, P., Aslan, A. and Cakici, A. (2009). Effect of air pollutant sulfur dioxide on phytohormone levels in some lichens. *Fres. Environ. Bull.*, 13(5), 436-440.
- Bedford, J.W., Roelofs, E.W. and M.J. Zabik. (1968). The freshwater mussel as a biological monitor of pesticide concentrations in a lotic environment. *Limnology and Oceanography*, 13(1), 118-126.
- Berberova, R., Bliznakov, A., Baykov, B. and Gyurov, R. (2008). Prinos za otshenka na okolnata sreda v oblast Stara Zagora. In: Proceeding of 7-th International Conference "Ecology-stable development", 23-25 October 2008, Vratsa, Bulgaria.
- Bianu, E. and Nica, D. (2006). Honeybees – bioindicators in a heavy metals polluted area. In: Proceedings of the Second European Conference of Apidology EurBee, Prague, Czech Republic, 10-16.
- Bilalov, F.S., Kolupaev, B.I., Kotov, Y.S., Musaramova, S.S. and Skrebneva, L.A. (1992). Bee products and environmental control. *Pchelovodstvo*, 12, 4-6.
- Bingol, A., Aslan, A. and Cakici. (2009). Biosorption of chromate anions from aqueous solution by a cationic surfactant-modified lichen (*Cladonia rangiformis* (L.)) *J. Hazard. Mater.*, 161, 747-752.
- Bogan, A.E. (1993). Freshwater bivalve extinctions (Mollusca: Unionoida): A search for causes. *American Zoologist*, 33, 599-609.
- Brodo, I.M. (1961). Transplant experiments with coricolous lichens using a new technique. *Ecology*, 42, 838-841.
- Browder, S. F., Johnson, D. H. and Ball, I. J. (2002). Assemblages of breeding birds as indicators of grassland condition. *Ecological Indicators*, 2, 257-270.
- Cairns, Jr. and Pratt, J.R. (1993). A history of biological monitoring using benthic macroinvertebrates. In: *Freshwater Biomonitoring and Benthic Macroinvertebrates*. Chapman & Hall, New York, Pp: 10-27.
- Caroline, J.L., Trevor, G.P., Andrew, A.M. and Kirk, T.S. (2001). Survival and behaviour of the earthworms *Lumbricus rubellus* and *Dendrodrilus rubidus* from arsenate-contaminated and non-contaminated sites *Soil Biology and Biochemistry*, 33, 1239-1244.
- Cavalchi, B. and Fornaciari, S. (1983). Api, miele, polline e propoli come possibili indicatori di un inquinamento da piombo e fluoro - Una esperienza di monitoraggio biologico nel comprensorio ceramico di Sassuolo-Scandiano.- In: *Atti del seminario di studi "i biologi e l'ambiente" Nuove esperienze per la sorveglianza ecologica*. Reggio Emilia, Italy (Manzini, P., Spaggiari, R., Eds) 17-18 febbraio, 275-300.
- Cicek, A., Koparal, A.S., Aslan, A. and Yazici, K. (2008). Accumulation of heavy metals from motor vehicles in transplanted lichens in an urban area. *Commun. Soil Sci. Plant Anal.*, 39, 168-176.
- Conti, M.E. and Cecchetti, G. (2001). Biological monitoring: lichens as bioindicators of air pollution assessment – a review. *Environmental Pollution*, 114, 471-492, Reprinted (or higher parts taken) with a kind permission from Elsevier.
- Coon, T.G., Eckblad, J.W. and P.W. Trygstad. (1977). Relative abundance and growth of mussels (Mollusca: Eulamellibranchia) in pools 8, 9, and 10 of the Mississippi River. *Freshwater Biology*, 7, 279-85.
- Cooperrider, A. and Noss, R. (1994). Saving aquatic biodiversity. *Wild Earth*. Spring, 54-64.
- Crane, E. (1984). Bees, honey and pollen as indicators of metals in the environment. *Bee World*, 65(1), 47 – 49.
- Crane, E. (1984). Bees, honey and pollen as indicators of metals in the environment. *Bee World*, 55, 47-49.
- Curtis, B., Roberts, K. S., Griffin, M., Bethune, S., Hay, C. J. and Kolberg, H. (1998). Species richness and conservation of Namibia freshwater macro-invertebrates, fish and amphibians. *Biodiversity and Conservation*, 7, 447-466.
- Duelli, P. and Obrist, M. K. (2003). Biodiversity indicators: the choice of values and measures. *Agriculture, Ecosystems and Environment*, 98, 87-98.
- Eguchi, S., Hatano, R. and Sakuma, T. (1995). Toshi effect of earthworms on the decomposition of soil organic matter. *Nippon Dojo-HiryogakuZasshi*, 66, 165-167.

- Ellis, M.M. (1942). Fresh-water impoundments. Transactions of the American Fisheries Society, 71st Annual Meeting, 80-93. Evermann, B.W., and H.W. Clark. 1918. The Unionidae of Lake Maxinkukee. Proceedings of the Indiana Academy of Science, 1917, 251-285.
- Eneva, S. and Todorova, M., 2004. Content of heavy metals in leached maroon forest soils in the region of the Agrobiochim – Stara Zagora. Journal of Agricultural Science and Forest Science, (2), 65-67.
- Fakhimzadeh, K., Lodenius, M., Kujala, J., Kahioloto, H. and Tulisalo, E. (2005). Using bees and hive products to assess metal pollution in Finland. In: Proceeding of XXXIX-th Apimondia International Apicultural congress, 21-26 August 2005, Dublin, Ireland, Pp: 52.
- Fenton, M.B. and Simmons, N.B. (2014). Bats: a world of science and mystery. University of Chicago Press, Chicago.
- Fu, C., Wu, J., Chen, J., Wu, Q. and Lei, G. (2003). Freshwater fish biodiversity in the Yangtze River basin of China: patterns, threats and conservation. Biodiversity and Conservation, 12, 1649–1685.
- Gallina, A., Baggio, A. and Mutinelli, F. (2005). Heavy metal contamination of honey in Veneto region (North-eastern Italy): an overview of the situation 2003 – 2004. In: bioindicators. Biological Reviews, 73, 181-201.
- Ghorbanil, M., Bakand, Z., Khaniki, B. and Bakand, S. (2007). Air pollution effects on the activity of antioxidant enzymes in Nerium oleander and Robinia pseudo acacia plants in Tehran. Iranian Journal of Environmental Health, Science and Engineering, 4(3), 157-162.
- Gupta, S., Bhattacharya, D., Datta, J.K., Nayek, S. and Satpati, S. (2009) Effects of vehicular emissions on biochemical constituents of leaves. Pollut. Res., 28, 157–160.
- Gupta, S., Nayek, S. and Bhattacharya, P. (2011) Effect of air-borne heavy metals on the biochemical signature of tree species in an industrial region, with an emphasis on anticipated performance index. Chemistry and Ecology, 27(4), 381-392.
- Haeba, M., Kuta, J., Arhouma, Z.K. and Elwerfalli, H.M.A. (2013). Earthworm as bioindicator of soil pollution around Benghazi City, Libya. J Environ Anal Toxicol., 3, 6.
- Hau, D.M., Chang, T.C., Huang, H.T. and Lin, W.D. (1976). Ecological studies on the stream water pollution near the water works in Taipei city. Report II. Analysis of biotic factors in the stream water. Biol. Bull. Natl. Taiwan Normal Univ., 11, 39-49.
- Hayes, A., Bakand, S. and Winder, C. (2007). Novel in vitro exposure techniques for toxicity testing and Biomonitoring of airborne contaminants. In: Drug Testing In vitro-Achievements and Trends in cell culture techniques, Wiley-VCH, Berlin, 103-124.
- Heino, J., Paavola, R., Virtanen, R. and Muotka, T. (2005). Searching for biodiversity indicators in running waters: do bryophytes, macroinvertebrates, and fish show congruent diversity patterns? Biodiversity and Conservation, 14, 415–428.
- Higginbottom, K., Green, R. and Northrope, C. (2003). A framework for managing the negative impacts of wildlife tourism on wildlife. Human Dimensions of Wildlife, 8, 1–24.
- Holt, E.A. and Miller, S.W. (2011). Bioindicators: using organisms to measure environmental impacts. Nature Education Knowledge, 3, 8. Honeybee biofilter abilities. In Proceeding of XXXIX-th Apimondia International Apicultural Congress, 21-26 August 2005, Dublin, Ireland, Pp: 95.
- Hong, C.C. (1979). The investigation of aquatic life in Tan-Shui river basin and the assessment of its pollution degree. Biol. Bull. Natl. Taiwan Normal Univ., 14, 23-31.
- Hughes, M.H. and Parmalee, P.W. (1999). Prehistoric and modern freshwater mussel (mollusca: bivalvia) faunas of the Tennessee River: Alabama, Kentucky, and Tennessee. Regulated Rivers: Research and Management, 15, 24-42.
- Inlay, M.J. (1982). The use of shells of freshwater mussels in monitoring heavy metals and environmental stresses: a review. Malacological Review, 15, 1-14.
- Jeliazkova, I., Marinova, M. and Gurgulova, K. (2002). Honey bee and their products as bioindicators for environmental pollution. Influence of sub-feeding of bee families with microelements contaminated sugar solution on content of those elements in bee organism. Journal of Animal Science, 39(4-5), 154-157.
- Jeliazkova, I., Marinova, M. and Peneva, V. (2001). Honey bee and their products as bioindicators for environmental pollution. Study on mineral content of body of bee-workers received different doses of microelements with their food. Journal of Animal Science, 38(6), 37-40
- Jones, G., 2012. What bioindicators are and why they are important. In: Flaquer, C., Puig-Montserrat, X. (Eds). Proceedings of the International Symposium on the Importance of Bats as Bioindicators. Museum of Natural Sciences Edicions, Granollers, Pp: 18-19.

- Jones, G., Jacobs, D.S., Kunz, T.H., Willig, M.R. and Racey, P.A. (2009). Carpe noctem: the importance of bats as bioindicators. *Endangered Species Research*, 8, 93-115.
- Joshi, N., Chauhan, A. and Joshi, P.C. (2009). Impact of industrial air pollutants on some biochemical parameters and yield in wheat and mustard plants. *Environmentalist*, 29, 398-404.
- Knopp, H., 1954. Ein neuer Weg zur Darstellung biologischer Gewässeruntersuchungen, erläutert an einem Gutellangsschnitt des Mains. *Die Wasserwirtschaft*, 45, 9-15.
- Kolkwitz, R. and Marrson, M. (1908). *Ökologie der pflanzlichen Saprobien*. *Ber. Deut. Bot. Ges.*, 26, 505-519.
- Landres, P.B., Verner, J. and Thomas, J.W. (1988). Ecological uses of vertebrate indicator species: a critique. *Conserv. Biol.*, 2, 316-328.
- Lee, C.D., Kuo, C.L. and Wang, S.P. (1967). Benthic macroinvertebrates and fish as biological indicator, with reference to the community diversity index. Report of Taiwan Prov. Inst. of Water Poll. Prot., 1, 1-44.
- Lindenmayer, D.B. and Fischer, J. (2003). Sound science or social hook: a response to Booker's application of the focal species approach. *Landsc. Urban Plann.*, 62, 149-158.
- Lindenmayer, D.B. and Likens, G.E. (2011). Direct measurement versus surrogate indicator species for evaluating environmental change and biodiversity loss. *Ecosystems*, 14, 47-59.
- Lindenmayer, D.B., Margules, C.R. and Botkin, D. (2000). Indicators of forest sustainability biodiversity: the selection of forest indicator species. *Conserv. Biol.*, 14, 941-950.
- Markert, B. (2008). From biomonitoring to integrated observation of the environment – the multi-markered bioindication concept. *Ecological Chemistry and Engineering*. 15(3), 315-330.
- Marking, L.L. and Bills, T.D. (1979). Acute effects of silt and sand sedimentation on freshwater mussels. Pp: 204-211 in Rasmussen, J.R., ed. *Proceedings of the UMRCC symposium on Upper Mississippi River bivalve mollusks*. Upper Mississippi River Conservation Committee, Rock Island, Illinois.
- McGeoch, M.A. (1998). The selection, testing and application of terrestrial insects as
- Merian, E., Anke, M., Ihnat, M. and Stoepler, M., 2008. *Elements and their compounds in the environment (Second edition)* published online. Copyright Wiley-VCH Verlag GmbH & Co KGaA.
- Morrison, M.L. (2009). *Restoring Wildlife: Ecological Concepts and Practical Applications*. Publication of Society for Ecological Restoration International. Island press, Washington.
- Mueller, P. (1980). *Biogeographie*. UTB, Ulmer Verlag. Stuttgart.
- Naimo, T.J. (1995). A review of the effects of heavy metals on freshwater mussels. *Ecotoxicology*, 4, 341-362.
- Neves, R.J. (1993). A state-of-the-union address. Conservation and management of freshwater mussels. Pp: 1-10.
- Newsome, D., Lewis, A. and Moncrieff, D. (2004). Impacts and risks associated with developing, but unsupervised, stingray tourism at Hamelin Bay, Western Australia. *International Journal of Tourism Research*, 6(5), 305-323.
- Ortmann, A.E. (1909). The destruction of the freshwater fauna in western Pennsylvania. *Proceedings of the American Philosophical Society*, 48(191), 90-110.
- Palacio, N.J., Verdu, J.R., Galante, E., Jimenez, G.D. and Olmos, O.G. (2007). Birds and fish as bioindicators of tourist disturbance in springs in semi-arid regions in Mexico: A basis for management. *Animal Biodiversity and Conservation*, 30, 1.
- Pandey, S.K., Tripathi, B.D., Prajapati, S.K., Mishra, V.K., Upadhyay, A.R., Rai, P.K. and Sharma, A.P. (2005). Magnetic properties of vehicle derived particulates and amelioration by *Ficus infectoria*: a keystone species. *Ambio: A Journal on Human Environment*, 34(8), 645-646.
- Park, K. (2015). Mitigating the impacts of agriculture on biodiversity: bats and their potential role as bioindicators. *Mammalian Biology*, this issue.
- Petkov, G., Yablanski, T., Todorova, M., Pavlov, D., Kostadinova, G. and Barakova, V. (2010). Ecological assessment of soils from parks and places in Stara Zagora. *Journal of Agricultural Science and Forest Science*, 9(4), 24-31.
- Porrini, C., Medrzycki, P., Bortolotti, L., Sabatini, A.G., Girotti, S., Ghini, S., Grillenzoni, F., Gattavecchia, E. and Celli, G. (2003). Honeybees as bioindicators of the environmental pollution. In: *Proceeding of XXXVIIIth Apimondia International Apicultural Congress*, 24-29 August 2003, Ljubljana, Slovenia, Final Programme and Book of Abstracts, Pp: 424.
- Posudin, Y. (2014). *Bioindication, in Methods of Measuring Environmental Parameters*. John Wiley

- & Sons, Inc., Hoboken, NJ, USA. Proceeding of XXXIX-th Apimondia International Apicultural congress, 21-26 August 2005, Dublin, Ireland, Pp. 145 – 146.
- Pyrovetsi, M. and Papastergiadou, A. (1992). Biological Biological conservation implications of water level fluctuations in wetlands of international importance: Lake Kerkini, Macedonia, Greece. *Environmental Conservation*, 19, 235–243.
- Raabe, O.G. (1999). Respiratory exposure to air pollutants. In: *Air pollutants and the respiratory tract*. Swift, D.L. & Foster, W.M. (Eds). Marcel Dekker INC, N. Y. USA., 39-73.
- Rossbach, M. and Lambrecht, S. (2006) Lichens as biomonitors: global, regional and local aspects. *Croatica Chemica Acta CCACAA*, 79(1), 119–124.
- Schuhmacher, M., Nadal, M. and Domingo, J.L. (2009). Environmental monitoring of PCDD/Fs and metals in the vicinity of a cement plant after using sewage sludge as a secondary fuel. *Chemosphere*, 74, 1502-1508.
- Sharma, A.P. and Tripathi, B.D. (2009). Biochemical response in tree foliage exposed to coal-fired power plant emission in seasonally dry tropical environment, *Environ. Monit. Assess*, 158, 197–212.
- Sharma, A.P., Rai, P.K. and Tripathi, B.D. (2007) Magnetic Biomonitoring of Roadside Tree Leaves as a Proxy of Vehicular Pollution. In: *Urban Planing and Environment: Strategies and Challenges*, Lakshmi Lakshmi Vyas (Ed.), Mc Millan Advanced Research Series, pp. 326-331.
- Showman, R.E. (1988). Mapping air quality with lichens, the North American experience. In: *Lichens, Bryophytes and air quality*. Nash, T.H., Writh, V. (Eds.). Cramer, Berlin, pp. 67-89.
- Simmons, G.M. and J.R. Reed. (1973). Mussels as indicators of biological zone recovery. *Water Pollution Control Federation*, 45(12), 2480-2492.
- Sladeczek, V., 1973. System of water quality from the biological point of view. *Ergebnisse. Der. Limnol.*, 7, 1-218.
- Spellerberg, I.F. (2005). *Monitoring Ecological Change*. Cambridge University Press, Cambridge.
- Spodniewska, A. and Romaniuk, K. (2006). Concentration of lead and cadmium in bees and bee bread. In: *Proceedings of the Second European Conference of Apidology EurBee*, Prague, Czech Republic, 10-16.
- Stanimirovic, Z., Stevanovic, J., Fakhimzadeh, K. and Baltic, M. (2005). Investigations of differences of heavy metals amount in honeybee bodies and their honey-indication.
- Svendsen, C., Spurgeon, D.J., Hankard, P.K. and Weeks, J.M. (2004). A review of lysosomal membrane stability measured by neutral red retention: is it a workable earthworm biomarker? *Ecotoxicology and Environmental Safety*, 57, 20-29.
- Svoboda, J. (1962). Teneur en strontium 90 dans les abeilles et dans leurs produits.- *Bulletin Apicole*, 5: 101-103.
- Takuchev, N. (2011). Morbidity from Non-Cancer Deceases, Associated with Air Pollution by Carbon oxide Of Stara Zagora, Bulgaria. In: *Proceeding of Food Science, Engineering and Technologies*, 14-15 October 2011, Plovdiv, Bulgaria, Pp: 223-226.
- Tershy, B. R., Breese, D. and Croll D. A. (1997). Human perturbations and conservation strategies for San Pedro Mártir Island, *Islas del Golfo de California Reserve, México*. *Environmental Conservation*, 24(3), 261–270.
- Wilkomirski, B. (2013). History of bioindication (*Historiabi bioindykacji*). *Monitoring Srodowiska Przyrodniczego*, 14, 137-142.
- Wurtz, C.B. (1956). Fresh-water mollusks and stream pollution. *Nautilus*, 69, 96-100.
- Yazici, K. and Aslan, A. (2006). Distribution of epiphytic lichens and air pollution in the city of Trabzon, Turkey. *Bull. Environ. Contam. Toxicol.*, 77, 838-845.
- Yeager, B.L. (1993). Dams. Pp: 57-113 in Bryan, C.F. and Rutherford, D.A eds. *Impacts on warmwater streams: guidelines for evaluation*. Warmwater Stream Committee, Southern Division, American Fisheries Society, Little Rock, Arkansas.
- Zelinka, M. and Marvan, P. (1961). Zur Prazisierung der biologischen klassifikation der Reinheit fliessender Gewasser. *Arch. Hydrobiol.*, 57, 389-407.
- Zhelyazkova, I., Marinova, M. and Gurgulova, K. (2004). Changes in the quantity of heavy metals in the haemolymph of worker bees fed micro-element contaminated sugar solution. *Aricilik*, 4(2), 77 – 80.
- Zukal, J., Pikula, J. and Bandouchova, H. (2015). Bats as bioindicators of heavy metal pollution: history and prospect. *Mammalian Biology*, this issue.

